

Cirrhosis in Cats

Overview

Cirrhosis is chronic, end-stage liver disease in which normal liver tissue has been replaced by fibrous scar tissue. Your cat needs approximately 20% of normal liver function to survive. When cirrhosis occurs, scar tissue replaces functioning liver cells. If normal liver function falls below 20%, the disease becomes terminal. Cirrhosis can happen at any age, but is most common in cats older than 7 years of age.

Cirrhosis occurs as a result of damage to the liver by many diseases, drugs, or toxins. Common diseases that may lead to cirrhosis include cancer as well as viral, bacterial, and [fungal infections](#) that cause hepatitis (inflammation of the liver). Certain toxins and long-term use of some medications, such as corticosteroids and common pain-relieving medications, can also cause cirrhosis. Therefore, it is crucial to have your cat's liver function monitored when he is on certain medications.

Symptoms

Symptoms will depend on the cause of the cirrhosis, and may include:

- [Loss of appetite](#); weight loss
- [Vomiting](#)
- [Diarrhea](#) or constipation
- Lack of energy; [depression](#)
- Increased thirst; urination
- Swollen abdomen (fluid-filled)
- Orange-tinged urine
- Yellow-tinged gums and whites of (sclera) or lining of the eyes ([jaundice](#))
- Bleeding problems
- Behavior changes; seizures; pacing or circling
- [Painful abdomen](#)
- Lack of coordination

Diagnosis/Treatment

Your veterinarian will take a complete history and perform a thorough physical exam of your pet. Additionally, diagnostic tests will be required to identify if your cat has cirrhosis of the liver. These may include:

- Chemistry tests to evaluate kidney, liver, and pancreatic function, as well as sugar levels
- Serologic tests to identify if your pet has been exposed to infectious diseases
- A complete blood count (CBC) to rule out certain

blood-related conditions

- Electrolyte tests to ensure your pet isn't dehydrated or suffering from an electrolyte imbalance
- Urine tests to screen for urinary tract infection and other disease
- A thyroid test to determine if the thyroid gland is functioning normally
- X-rays to evaluate the size, shape, and position of the liver
- An abdominal ultrasound to evaluate the liver and other vital organs
- Coagulation profiles to assess your pet's clotting function
- A liver biopsy

Treatment for your cat will vary depending on the underlying cause of the liver damage and cirrhosis. The good news is that treating the underlying cause of cirrhosis can, in many cases, stop the progression of the damage.

Treatment may include the following:

- Discontinuation of any therapy that may have caused the liver damage
- Intravenous fluid and electrolyte therapy, if your pet is dehydrated
- Blood products, if your pet is anemic
- Dietary modifications
- Medications, depending on the cause

Prevention

The prognosis for your cat depends on two factors: how much of the liver's function is impaired and the ability to treat and control the underlying cause. The most effective prevention of cirrhosis is to address liver disease as early as possible and maintain your cat's preventive health care to avoid any situation that may cause liver disease.

If you have any questions or concerns, you should always visit or call your veterinarian – they are your best resource to ensure the health and well-being of your pets.